

# STAT 302: Assignment 1 Solutions

Summer 2026, Due May 20th, 11:59pm on Gradescope.

## Question 1 [6 points]

Suppose that a student's phone screentime report shows that they are scrolling Reddit 50% of days, checking Instagram 80% of days, and checking their email 10% of days. 30% of the days, the student checks Reddit and Instagram in the same day (we don't know whether or not they also check their email on these days). 17% of the days, they exclusively look at Reddit. They never check their email without opening Reddit or Instagram. 3% of the days they check all three apps.


Show your work and all calculations. You may use visuals to help you.

- (a) On a randomly selected day, what is the probability that the student checks only Reddit and their email?
- (b) On a randomly selected day, what is the probability that the student only checks Instagram, but not email nor Reddit?
- (c) What is the probability that they do not check any of the three apps?


## Solution

- (a) 3%
- (b) 46%
- (c) 0%


You can fill out a Venn Diagram to easily solve for each piece:




(a) Step 1




(a) Step 2




(a) Step 3



(a) Step 4



(a) Step 5



Total = 100. No days without an app.

(a) Step 6

## Question 2 [8 points]

Prove the Bonferroni inequality:

$$\mathbb{P}\left(\bigcap_{i=1}^n A_i\right) \geq \sum_{i=1}^n \mathbb{P}(A_i) - (n-1)$$

Show every step of the proof. List any results, rules, or theorems you use in the respective lines of your proof.

Hint:

$$\bigcap_{i=1}^n A_i = \left(\bigcup_{i=1}^n A_i^c\right)^c.$$

### Solution

$$\begin{aligned} \mathbb{P}\left(\bigcap_{i=1}^n A_i\right) &= \mathbb{P}\left(\left(\bigcup_{i=1}^n A_i^c\right)^c\right) \\ &= 1 - \mathbb{P}\left(\bigcup_{i=1}^n A_i^c\right) && \text{complement rule} \\ &\geq 1 - \sum_{i=1}^n \mathbb{P}(A_i^c) && \text{Boole's inequality} \\ &= 1 - \sum_{i=1}^n (1 - \mathbb{P}(A_i)) && \text{complement rule} \\ &= 1 - \sum_{i=1}^n 1 + \sum_{i=1}^n \mathbb{P}(A_i) \\ &= 1 - n + \sum_{i=1}^n \mathbb{P}(A_i) \\ &= \sum_{i=1}^n \mathbb{P}(A_i) - (n-1). \end{aligned}$$

## Question 3 [7 points]

Prove that  $\mathbb{P}(\cdot | B)$  is a probability. Hint: there are three axioms in the formal definition of a probability.

### Solution

We must show from the definition that this function satisfies the 3 axioms.

**Axiom 1:**  $\mathbb{Q}_B(\Omega) = 1$ :

$$\mathbb{Q}_B(\Omega) = \frac{\mathbb{P}(\Omega \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B)}{\mathbb{P}(B)} = 1$$

**Axiom 2:**  $\mathbb{Q}_B(A) \geq 0$  for any  $A \in \Omega$ :

$$\mathbb{Q}_B(A) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} \geq 0$$

**Axiom 3:** If  $\{A_i\}_{i \geq 1}$  are disjoint events:

$$\begin{aligned} \mathbb{Q}_B\left(\bigcup_{i=1}^{\infty} A_i\right) &= \mathbb{P}\left(\left[\bigcup_{i=1}^{\infty} A_i\right] \cap B\right) / \mathbb{P}(B) \\ &= \mathbb{P}\left(\bigcup_{i=1}^{\infty} [A_i \cap B]\right) / \mathbb{P}(B) \\ &= \left[ \sum_{i=1}^{\infty} \mathbb{P}(A_i \cap B) \right] / \mathbb{P}(B) \\ &= \sum_{i=1}^{\infty} \frac{\mathbb{P}(A_i \cap B)}{\mathbb{P}(B)} \\ &= \sum_{i=1}^{\infty} \mathbb{Q}_B(A_i) \end{aligned}$$

### Question 4 [8 points]

A quality control inspector examines items coming off a production line and classifies each as defective (D) or non-defective (N). The inspector examines items one at a time and **stops as soon as they find 2 defective items, or after examining 4 items**, whichever comes first.

- Write out the sample space  $\Omega$ .
- Let  $A$  be the event that the inspector stops after exactly 3 items. List the outcomes in  $A$ .
- Let  $B$  be the event that fewer than 2 defective items are found in total. List the outcomes in  $B$ .
- What is  $A \cap B$ ? Describe this subset in words.

### Solution

(a) [ 2 points]

$$\Omega = \{DD, NDD, DND, NNDD, NDND, DNND, NNNN, DNNN, NDNN, NNDN, NNND\}$$

(b) [2 points]

If the inspector stops after three items, it means they found two defects, and these two defects were not the first two.  $A = \{NDD, DND\}$ .

(c) [2 points]

If there are fewer than two defective items in total, then that means there are either 0 or 1 defectives across the whole sequence. And that means we'd need four items.

$$B = \{NNNN, DNNN, NDNN, NNDN, NNND\}$$

(d) [2 points]

Sequences where the inspector stopped after three items and there were fewer than 2 defects. This isn't possible as an inspector will only stop at 3 items if there are 2 defects. Therefore,  $A \cap B = \emptyset$